

**CONTENT**

	<b>Page</b>
● Frequency distribution and potence ratio of agronomic traits in F1 generation of intraspecific crosses of Robusta coffee. <b>Ari Wibowo and Ucu Sumirat</b> .....	83-97
● Adaptation of introduced Robusta coffee clones in some agroclimate types in East Java. <b>Pingkan Sugianto, Sudarsono, Dewi Sukma, and Ucu Sumirat</b> .....	98-107
● Establishment of an efficient primary callus induction for embryogenic potential of <i>Coffea arabica</i> L. <b>Rina Arimarsetiowati, Budi Setiadi Daryono, Yohana Theresia Maria Astuti, and Endang Semiarti</b> .....	108-119
● Flush characteristics of several cocoa genotypes different in resistant to vascular streak dieback. <b>Indah Anita-Sari, Bayu Setyawan, Agung Wahyu Susilo, Nurhadini Fitri Isnaini, Samsul Paputpungan, Febrilia Nur'aini, and Nur Solecha Ruseani</b> .....	120-127
● Caffeine content of Bondowoso Arabica ground coffee with variation of roasting profile and type of packages. <b>Dyah Ayu Savitri, Ayu Puspita Arum, Hasbi Mubarak Suud, Oria Alit Farisi, Susan Barbara Patricia SM, Bambang Kusmanadhi, and Denna Eriani Munandar</b> .....	128-137
● Cocoa fresh beans aqueous extract as free radical quencher and ferric reducer. <b>Samuel Yap Kian Chee, Arief Huzaimi Bin M.D. Yusof, and Hui Han Chin</b> .....	138-144
● Covid-19 pandemic awareness and coping strategies of cocoa farmers in Ondo State of Nigeria. <b>Foluso Temitope Agulanna</b> .....	145-154

